

Janet T. Mills  
Governor

Jeanne M. Lambrew, Ph.D.  
Commissioner



Maine Department of Health and Human Services  
Maine Center for Disease Control and Prevention  
11 State House Station  
286 Water Street  
Augusta, Maine 04333-0011  
Tel; (800) 821-5821; Fax (207) 287-6865  
TTY: Dial 711 (Maine Relay)


## Maine Monthly Pertussis Surveillance Report

### September 2019

Date of Report: Oct 16, 2019

This update summarizes cases of pertussis reported to the Maine Center for Disease Control from January 1<sup>st</sup> through September 30<sup>th</sup>, 2019. There were 15 pertussis cases reported statewide in September, compared to 43 reported cases in 2018 and a 5-year median of 20 during that same time (see below graph). Six counties had rates of pertussis above the state rate: Cumberland, Hancock, Knox, Lincoln, Waldo and Washington. Year to date, most cases were in kids between the ages of 11-19 (50.8%), the average age was 15.1 years and the median age was 13 years.

No outbreaks were reported in September.



- The clinical case definition and surveillance case definitions of pertussis are different, so the reported number of cases may seem lower than expected
- All data presented is preliminary and subject to change

Janet T. Mills  
Governor



Jeanne M. Lambrew, Ph.D.  
Commissioner

Maine Department of Health and Human Services  
Maine Center for Disease Control and Prevention  
11 State House Station  
286 Water Street  
Augusta, Maine 04333-0011  
Tel; (800) 821-5821; Fax (207) 287-6865  
TTY: Dial 711 (Maine Relay)

**Table 1. 2019 Monthly Case Count, YTD Case Count and YTD Incidence Rate per 100,000**

County	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total Cases (YTD)	Incidence Per 100,000 (YTD)
Aroostook County	0	0	0	0	0	0	0	0	0				0	0.0
Androscoggin County	4	1	3	0	0	2	0	0	1				11	10.3
Cumberland County	3	4	17	11	13	18	7	7	1				81	27.9
Franklin County	0	0	0	0	0	0	0	0	0				0	0.0
Hancock County	16	9	5	4	8	4	6	9	4				65	118.9
Kennebec County	0	0	2	1	2	0	0	0	0				5	4.2
Knox County	2	4	1	1	8	8	3	3	1				31	77.8
Lincoln	2	4	2	1	0	1	1	1	0				12	35.3
Oxford County	0	1	0	1	1	0	1	4	1				9	15.7
Penobscot County	1	0	2	0	0	0	0	2	2				7	4.6
Piscataquis County	0	0	0	0	0	0	0	1	0				1	5.9
Sagadahoc County	1	0	0	0	0	0	0	0	0				1	2.8
Somerset County	0	1	0	0	0	0	0	0	1				2	3.9
Waldo County	10	11	4	5	3	1	1	3	0				38	97.1
Washington County	3	2	2	3	0	0	0	3	0				13	41.1
York County	13	3	3	6	4	2	3	3	4				41	20.4
<b>Maine</b>	<b>55</b>	<b>40</b>	<b>41</b>	<b>33</b>	<b>39</b>	<b>36</b>	<b>22</b>	<b>36</b>	<b>15</b>				<b>317</b>	<b>23.8</b>

**Table 2. Pertussis cases by age group- Maine, Year to Date 2019**

Age Group	# of Cases	Percent of Cases
< 6 Months	5	1.5%
6-11 Months	12	3.6%
1-6 Years	60	18.2%
7-10 Years	39	11.9%
11-19 Years	167	50.8%
≥ 20 Years	46	14.0%

**Demographics: Average age: 15.1 years,  
Median age: 13 years. Min 1 month, Max 87  
years**

**51% Female, 49% Male**

- The clinical case definition and surveillance case definitions of pertussis are different, so the reported number of cases may seem lower than expected
- All data presented is preliminary and subject to change

Janet T. Mills  
Governor

Jeanne M. Lambrew, Ph.D.  
Commissioner



Maine Department of Health and Human Services  
Maine Center for Disease Control and Prevention  
11 State House Station  
286 Water Street  
Augusta, Maine 04333-0011  
Tel; (800) 821-5821; Fax (207) 287-6865  
TTY: Dial 711 (Maine Relay)

Table 3. Pertussis Outbreaks by County and Facility Type- Maine, Year to Date 2019

County	Facility Type		
	School (K-12)	Daycare	Other
Cumberland	4		
Hancock	2		
Lincoln	2		
Washington			1
York	2		

### Pertussis Prevention:

- The best way to prevent pertussis (whooping cough) among babies, children, teens, and adults is to get vaccinated.
  - For more information on pertussis vaccines visit <https://www.cdc.gov/pertussis/vaccines.html>
- Pertussis spreads by coughing and sneezing while in close contact with others, who then breathe in the bacteria. Practice good hand hygiene to prevent the spread of respiratory illnesses. To practice good hygiene, you should:
  - Cover your mouth and nose with a tissue when you cough or sneeze.
  - Put your used tissue in the waste basket.
  - Cough or sneeze into your upper sleeve or elbow, not your hands, if you don't have a tissue.
  - Wash your hands often with soap and water for at least 20 seconds or use an alcohol-based hand rub if soap and water are not available.

- The clinical case definition and surveillance case definitions of pertussis are different, so the reported number of cases may seem lower than expected
- All data presented is preliminary and subject to change